

A Comparison Of Proximal Femoral Nail Antirotation And

Yeah, reviewing a book **a comparison of proximal femoral nail antirotation and** could go to your close links listings. This is just one of the solutions for you to be successful. As understood, feat does not suggest that you have astounding points.

Comprehending as without difficulty as concurrence even more than new will have enough money each success. neighboring to, the publication as without difficulty as perception of this a comparison of proximal femoral nail antirotation and can be taken as without difficulty as picked to act.

domain Public Library provides a variety of services available both in the Library and online. ... There are also book-related puzzles and games to play.

A Comparison Of Proximal Femoral

A Comparison of the Proximal Femoral Nail Antirotation Device and Dynamic Hip Screw in the Treatment of Unstable Peritrochanteric Fracture Show all authors. YZ Xu * YZ Xu. See all articles by this author. Search Google Scholar for this author, DC Geng * DC Geng. See all articles by this author.

A Comparison of the Proximal Femoral Nail Antirotation ...

Objective: To compare the effectiveness of proximal femoral nail antirotation (PFNA) and reconstruction nail with minimally invasive technique for ipsilateral femoral shaft and extracapsular hip fractures in young and middle-aged patients. Methods: Sixty-nine young and middle-aged patients with ipsilateral femoral shaft and extracapsular hip fractures were treated between January 2000 and ...

[Comparison of proximal femoral nail antirotation and ...

Free Online Library: Comparison of Proximal Femoral Geometry and Risk Factors between Femoral Neck Fractures and Femoral Intertrochanteric Fractures in an Elderly Chinese Population.(Original Article, Report) by "Chinese Medical Journal"; Health, general Aged Health aspects Care and treatment Comparative analysis Risk factors Fractures (injuries) Implants, Artificial Usage Medical records ...

Comparison of Proximal Femoral Geometry and Risk Factors ...

A comparison of proximal femoral locking compression plates with dynamic hip screws in extracapsular femoral fractures Author links open overlay panel B. Zhong a 1 Y. Zhang b 1 C. Zhang a C.-F. Luo a Show more

A comparison of proximal femoral locking compression ...

Objective: To compare the effects of proximal femoral nail antirotation blade (PFNA) and reverse less invasive stabilization system-distal femur (Liss-DF) systems in the treatment of proximal femoral fractures. Methods: Between June 2007 and October 2009, 41 proximal femoral fractures were treated, 22 with PFNA (group A) and 19 with reverse LISS-DF plates (group B).

Comparison of proximal femoral nail antirotation blade and ...

A comparison of the proximal femoral nail antirotation device and dynamic hip screw in the treatment of unstable peritrochanteric fracture. Xu YZ(1), Geng DC, Mao HQ, Zhu XS, Yang HL. Author information: (1)Department of Orthopaedics, First Affiliated Hospital of Soochow University, Suzhou, China. xuyaozeng@163.com

A comparison of the proximal femoral nail antirotation ...

Uzer et al. Comparison of two types of proximal femoral nails in the treatment of intertrochanteric femur fractures data were collected from charts and the outcomes of the most recent postoperative follow-up visits. Inclusion and Exclusion Criteria Patients ≥60 years of age at the time of surgery were in-cluded into the study.

Comparison of two types of proximal femoral nails in the ...

To compare the efficacy of three-point locating versus routine locating techniques for implanting helical blades for proximal femoral nail anti-rotation-II in the treatment of trochanteric fractures. From January 2010 to June 2013, 90 patients with intertrochanteric fractures were surgically treated, including 48 males and 42 females with an average age of 70.5 ± 7.2 years.

Comparison of two methods of locating proximal femoral ...

connecting line of the femoral condyle and the horizontal line was measured at the distal end of the femur, and the angle between the femoral neck and the horizontal plane was measured at the proximal end of the femur. The dif-ference between these two angles is the femoral neck ante-version (FNA), which is 9.3+6.68=15.98° (Fig. 1d, e).

Comparison of two methods of locating proximal femoral ...

Purpose. There is no consensus about the best option of internal fixation for unstable intertrochanteric fractures. The aim of the present study was to compare proximal femoral nail (PFN) with contralateral reverse distal femoral locking compression plate (reverse-DFLCP) in the management of unstable intertrochanteric fractures with compromised lateral wall.

Proximal femoral nails compared with reverse distal ...

Proximal Femoral Nail in Proximal Femoral Fractures – A Prospective Comparative Study, J Res Med Dent Sci, 2017, 5 (2):1-9 , DOI: 10.5455/jrmds.2017.521 Corresponding author : Shoab Shaikh

[PDF] Comparison of Intertan Nail Vs Proximal Femoral Nail ...

ing proximal femoral fracture remains a matter of debate. The aim of this study was to compare the clinical out-comes of proximal femoral fractures after PFNA and Liss-DF fixation. Materials and Methods Patients' data Between June 2007 and October 2009, 41 patients with fresh, closed proximal femoral fractures were admitted to our hospital.

Comparison of proximal femoral nail antirotation blade and ...

Abstract. The goal of this study was to compare the results of proximal femoral nail antirotation (PFNA; Synthes, Oberdorf, Switzerland) fixation with those of Gamma nail 3 (GN-3; Stryker, Mahwah ...

Treatment of Unstable Proximal Femoral Fractures ...

These results can be used to increase the awareness of proximal hip fracture among physicians and improve treatments and outcomes. To compare the BMD of the femoral neck region and the intertrochanteric region between the femoral neck fracture group and the intertrochanteric fracture group.

Comparison of hip subregion bone mineral density to the ...

PurposeThe aim of this study was to compare the results of intramedullary fixation with those of plate-screw fixation for peritrochanteric femoral fracture patients older than 60 years old.MethodsThis article reports on a retrospective review of patients who had peritrochanteric femoral fractures and were treated with a 95° fixed-angle screw plate (DCS) or an intramedullary nailing system (PFNA).

[PDF] Comparison of proximal femoral nail antirotation ...

Objectives . This biomechanical study was conducted to compare fixation stability of the proximal fragments and their mechanical characteristics in proximal femur models of unstable basicervical IT fractures fixed by cephalomedullary nailing using 3 different types of the femoral head fixation. Methods . A total of 36 composite femurs corresponding to osteoporotic human bone were used.

Biomechanical Comparison of Three Different Intramedullary ...

The goal of this study was to compare the results of proximal femoral nail antirotation (PFNA; Synthes, Oberdorf, Switzerland) fixation with those of Gamma nail 3 (GN-3; Stryker, Mahwah, New Jersey) fixation for unstable proximal femoral fractures. A total of 136 unstable trochanteric fractures were divided into 2 treatment groups: 66 patients were treated with the PFNA and 70 were treated with the GN-3.

Treatment of unstable proximal femoral fractures ...

The dynamization of the femoral neck screw or femoral neck blade through the nail was determined by measuring the distance between the most proximal point of the femoral neck screw/femoral neck ...

The Dynamics of Proximal Femoral Nails: A Clinical ...

The lateral femoral cutaneous nerve (LFCN) divides into several branches innervating the lateral and anterior aspects of the thigh.Indications are postoperative analgesia for hip surgery, meralgia paresthetica, and muscle biopsy of the proximal lateral thigh.